

Citation for published version:

Roberts, T, Daniels, J, Hulme, W, Hirst, R, Horner, D, Lyttle, MD, Samuel, K, Graham, B, Reynard, C, Barrett, M, Foley, J, Cronin, J, Umana, E, Vinagre, J & Carlton, E 2021, 'The COVID-19 emergency response assessment study: a prospective longitudinal survey of frontline doctors in the UK and Ireland', *BMJ Open*, vol. 11, no. 7, e049680. <https://doi.org/10.1136/bmjopen-2021-049680>

DOI:

[10.1136/bmjopen-2021-049680](https://doi.org/10.1136/bmjopen-2021-049680)

Publication date:

2021

Document Version

Peer reviewed version

[Link to publication](#)

Publisher Rights

CC BY-NC

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Title Page

Psychological Distress and Trauma in Doctors Providing Frontline Care During the COVID-19 Pandemic in the United Kingdom and Ireland: A Prospective Longitudinal Survey Cohort Study

Authors: Tom Roberts^{1, 2}, Jo Daniels³, William Hulme⁴, Robert Hirst⁵, Daniel Horner^{1, 6}, Mark D Lyttle^{2, 7}, Katie Samuel⁵, Blair Graham^{8, 9}, Charles Reynard¹⁰, Michael J Barrett^{11, 12}, James Foley¹³, John Cronin¹⁴, Etimbuk Umana¹⁵, Joao Vinagre¹⁶ and Edward Carlton^{1, 17} on behalf of The Trainee Emergency Research Network (TERN), Paediatric Emergency Research in the UK and Ireland (PERUKI), Research and Audit Federation of Trainees (RAFT), Irish Trainee Emergency Research Network (ITERN and Trainee Research in Intensive Care (TRIC))

Affiliations:

- 1) The Royal College of Emergency Medicine, London, UK
- 2) Emergency Department, Bristol Royal Hospital for Children, UK
- 3) Department of Psychology, University of Bath, UK
- 4) Statistical Consultant, Oxford, UK
- 5) Department of Anaesthesia, North Bristol NHS Trust, UK
- 6) Emergency Department, Salford Royal Hospital NHS Foundation Trust
- 7) Faculty of Health and Applied Sciences, University of the West of England, Bristol
- 8) Faculty of Health, University of Plymouth, Plymouth, UK.
- 9) Emergency Department, University Hospitals Plymouth, UK
- 10) Department of Cardiovascular Sciences, University of Manchester
- 11) Department of Emergency Medicine, Children's Health Ireland at Crumlin, Ireland
- 12) Women's and Children's Health, School of Medicine, University College Dublin, Ireland
- 13) Emergency Department, University Hospital Waterford, Waterford, Ireland
- 14) Department of Emergency Medicine, St Vincent's University Hospital, Dublin, Ireland
- 15) Emergency Department, Connolly Hospital Blanchardstown, Dublin, Ireland.
- 16) College of Anaesthesiologists of Ireland, Dublin, Ireland
- 17) Emergency Department, North Bristol NHS Trust

Corresponding Author

Dr Tom Roberts

Address: 12 Hamilton Road, Bristol, BS3 1PB

Email: Tomkieranroberts@gmail.com

Telephone: 07894234121

Abstract Word Count: 304

Manuscript Word Count: 3561

Objectives

The psychological impact of the COVID-19 pandemic on doctors is a significant concern. Due to the emergence of multiple pandemic waves, longitudinal data on the impact of COVID-19 is vital to ensure an adequate psychological care response. The primary aim was to assess the prevalence and degree of psychological distress and trauma in frontline doctors during the acceleration, peak and deceleration of the COVID-19 first wave. Personal and professional factors associated with psychological distress are also reported.

Design

A prospective online three-part longitudinal survey.

Setting

Acute hospitals in the UK and Ireland.

Participants

Frontline doctors working in Emergency Medicine (EM), Anaesthetics and Intensive Care Medicine (ICM) during the first wave of the COVID-19 pandemic in March 2020.

Primary outcome measures

Psychological distress and trauma measured using the General Health Questionnaire-12 and the Impact of Events-Revised.

Results

The initial acceleration survey distributed across networks generated a sample of 5440 doctors. Peak and deceleration response rates from the original sample were 71·6% (n=3896) and 56·6% (n=3079) respectively. Prevalence of psychological distress was 44·7% (n=1334) during the acceleration, 36·9% (n=1098) at peak and 31·5% (n=918) at the deceleration phase. The prevalence of trauma was 23·7% (n=647) at peak and 17·7% (n=484) at deceleration. The prevalence of probable post-traumatic stress disorder was 12·6% (n=343) at peak and 10·1% (n=276) at deceleration. Worry of family infection due to clinical work was the factor most strongly associated with both distress ($R^2 = 0\cdot06$) and trauma ($R^2 = 0\cdot10$).

Conclusion

Findings reflect a pattern of elevated distress at acceleration and peak, with some natural recovery. It is essential that policymakers seek to prevent future adverse effects through (a) provision of vital equipment to mitigate physical and psychological harm (b) increased awareness and recognition of signs of psychological distress and (c) the development of clear pathways to effective psychological care.

Trial Registration: ISRCTN 10666798

Strength and limitations of this study

- This paper presents key findings from a large cross-sectional longitudinal survey of practising emergency, anaesthetic and intensive care doctors in UK and Ireland during the acceleration, peak and deceleration of the first wave of the COVID-19 pandemic.
- This study provides an insight into the associated personal and professional factors associated with trauma and distress and could be utilised to identify those doctors who will most benefit from psychological interventions.
- Variation in regional peaks may have influenced accurate capturing of psychological distress and trauma rates and have not been accounted for.
- The findings cannot be extrapolated to longer-term psychological impact, and future work is planned to capture this.

Introduction

Clinicians providing frontline care have become central to the primary reception, assessment, and ongoing hospital treatment of patients with suspected Coronavirus Infectious Disease 2019 (COVID-19). These include doctors working in Emergency Medicine (EM), Anaesthetics and Intensive Care Medicine (ICM). Whilst this healthcare workforce is highly resilient and accustomed to facing traumatic situations, the COVID-19 pandemic has imposed unprecedented demands in workload intensity and personal health risk.^{1–4} High infection rates have been reported in frontline clinicians, with over 150 fatalities in the UK by May 2020.⁵ These factors are likely to affect psychological wellbeing, increasing the risk of traumatic stress both in the acute phase of the pandemic and at long-term follow up.^{6–9} Exposure to infectious disease outbreaks and elevated psychological distress have previously been associated with increased sickness rates, absenteeism, impaired performance at work, and the development of physical health problems.^{10–12} There is also an emerging evidence base from around the world of the psychological impact on healthcare workers.^{13–16} During the current COVID-19 pandemic there has been a global media focus on health and care workers with widespread public support.¹⁷ However, there is increasing recognition amongst key opinion leaders and psychological societies that this pandemic will lead to an unparalleled, though as yet unquantified, impact upon the psychological wellbeing of healthcare workers.^{18,19}

Studies evaluating psychological wellbeing in frontline clinicians during infectious disease outbreaks (including COVID-19) have demonstrated negative impacts may be significant.^{10,20,21} Systematic reviews and meta-analyses converge around common predictors of psychological distress following traumatic events, many of which are relevant to frontline clinicians. Key factors include preparedness, training, social and occupational support, exposure and threat to life, media use and history of mental health problems.^{1,7,21–23} However, these data have largely been collected as a snapshot either during or following outbreaks or as cross-sectional surveys in highly selected or self-selecting cohorts. Longitudinal data which describe evolving and cumulative effects on the psychological wellbeing of frontline working during the COVID-19 pandemic are therefore urgently required. Such studies are essential to understand and mitigate psychological impacts of future events upon this vital workforce and inform the development of policy and interventions.

The primary aim of this study was to assess the prevalence and degree of psychological distress and trauma in doctors providing frontline care during the acceleration, peak and deceleration phases of the COVID-19 pandemic. We also sought to establish which personal and professional factors were significantly associated with psychological distress at these time points.

Methods

Study Design and Participants

The “COVID-19 Emergency Response Assessment (CERA) Study” was a prospective online longitudinal survey of frontline doctors across the UK and Ireland undertaken during the acceleration, peak and deceleration phases of the first COVID-19 pandemic wave.²⁴ Doctors of all grades working in EM, Anaesthetics or ICM during the acceleration phase were invited to participate.

Procedures

This survey study is reported in line with Checklist for Reporting Results of Internet E-surveys (CHERRIES) guidelines.²⁵ Full details of survey distribution, design, administration, and time-points are available in the published protocol.²⁴ In brief, the survey was initially distributed during the acceleration phase of the first pandemic wave through research networks, training faculties or Royal College Networks via email or instant messaging groups, coordinated by identified site/region leads. The participation link was not shared on wider social media platforms, to avoid international contamination. At completion of the acceleration phase survey, participants entered personal email addresses for direct approach at peak and deceleration phases with a unique survey link to avoid duplication. The study was registered at the ISRCTN (10666798).

The acceleration, peak and deceleration surveys were developed iteratively by the study team and underpinned by evidence, or by consensus where necessary. Psychometric tools were selected by consensus of the study team, considering validity and utility of a range of standardised measures, balanced against the feasibility of delivery and completion by individuals likely to be working at maximum capacity.

Study data were collected and managed using REDCap (Research Electronic Data Capture) hosted at University Hospitals Bristol and Weston NHS Foundation Trust.^{26,27} Acceleration, peak and deceleration phases were defined a priori and adapted from the United States Centre for Disease Control “Preparedness and Response Frameworks for Influenza Pandemics”.²⁸ For each survey, exact survey distribution dates were decided per protocol by team consensus according to available public health data on number of confirmed cases (acceleration phase; UK: 18/03/2020 – 26/03/2020, Ireland: 25/03/2020 – 02/04/2020), nationally available COVID-19 daily death rates (peak phase; UK: 21/04/2020 – 05/05/2020, Ireland: 28/04/2020 – 12/05/2020) and at 30 days after distribution of the peak phase survey (deceleration phase; UK: 03/06/2020 – 17/06/2020, Ireland: 10/06/2020 – 24/06/2020). Ethical approval was obtained from the University of Bath (UK) and Children’s Health Ethics Committee (Ireland). Regulatory approval was obtained from the Health Regulation Authority (UK), Health and Care Research Wales. Participants provided electronic informed consent for each survey.

Survey Questions

Survey questions collected data for both the primary and secondary outcomes. Items included the General Health Questionnaire-12 (GHQ-12; provided with licence fee waived by GL Assessments, London, UK) for distress, and the Impact of Events Scale- Revised (IES-R; off licence) for trauma. Personal and professional characteristics relating to participants' current role, and their preparedness and experiences during the pandemic were collected. These were used as secondary outcome measures and are provided in full in the protocol and online supplement.²⁴

Outcomes

There were two co-primary outcomes in this survey: psychological distress, and trauma, as defined by the GHQ-12 and the IES-R respectively.

Distress – GHQ-12

The GHQ-12 is a 12-item self-report measure devised to screen for psychological distress in the general population.²⁹ The measure has high specificity and sensitivity, with reliability demonstrated across a range of populations.^{30,31} The GHQ-12 has been used in similar clinician-based studies measuring the psychological impact of infectious outbreaks and was chosen due to the brevity of the measure and its suitability for time-pressured medical staff.²¹ The GHQ-12 assesses current state and asks the participants to compare to usual state. GHQ-12 was asked at all 3 survey phases. Case level distress is defined as a score of more than 3.³⁰

Trauma – IES-R

The IES-R is a 22-item measure commonly used to measure post-traumatic stress following a pre-specified traumatic incident and has been used to evaluate the impact of infectious disease outbreaks on hospital staff.^{21,32} It contains 8 items that focus on 'intrusion', 8 items on 'avoidance' and 6 items on 'hyperarousal'. The IES-R was used at the peak and deceleration survey phases. A score of 24 or above indicates a clinically significant traumatic stress response, a score above 33 indicates best cut-off for a diagnosis of 'probable post-traumatic stress disorder' (PTSD).^{33,34}

The secondary outcomes captured included personal and professional characteristics and their association with psychological distress and trauma. These personal and professional factors were identified through rapid literature review of high-quality systematic reviews and meta-analysis by experts in pandemic research.^{1,21–23} All factors identified as predictors of outcome were retained. This was supplemented by factors deemed of specific or emerging interest by the expert study steering committee. These were defined a priori in the study protocol, with the exception of ethnicity which was added during the peak survey due to the specific emergence of ethnicity as a potential marker of poor physical health outcomes.²⁴

Statistical Analysis

The statistical analysis is described in detail in the published protocol.²⁴ GHQ-12 items were reported using two methods. In the first method, item responses are assigned to the values 0, 0, 1, 1 (from the

most positive to the most negative sentiment) and summed to form an aggregate score from zero (least distressed) to 12 (most distressed). Using this method, a score of more than 3 is indicative of case-level distress.³⁰ The second method assigns responses to 0, 1, 2, 3 (positive to negative sentiment) producing a score in the range 0 to 36, with zero representing the most healthy response (no psychological distress) and 36 the most unhealthy (maximal psychological distress). By presenting the two different scoring methods, we can both report the prevalence of case level distress across the sample (0-0-1-1 scoring method) and more sensitively detect changes within the sample over the three phases of the pandemic (0-1-2-3 scoring method).

IES-R responses were analysed by assigning the responses to 0, 1, 2, 3, 4 (positive to negative) producing a score in the range 0 (no trauma) to 88 (maximal trauma). A score of 24 or above indicates a clinically significant traumatic stress response, a score above 33 indicates best cut-off for a diagnosis of 'probable post-traumatic stress disorder' (PTSD).^{33,34}

The change over time in the GHQ-12 (phases 1, 2, and 3) and IES-R scores (phases 2 and 3) amongst participants who responded to all three surveys was examined with repeated measures linear mixed-effect models, with survey phase as the single fixed effect and a participant-level random effect. These model describe the association between pandemic phase and psychological distress (GHQ-12) and trauma (IES-R).

To identify potential modifiers of the change in GHQ-12-score or IES-R-score over time, further models were constructed for each of the measured personal and professional variables. Each model included the single variable of interest, survey phase, their interaction (to allow for a change in the association between the outcome and the variable over time), and a participant-level random effect as before.. Responses where the variable value was missing were removed.³⁵ Nagakawa's marginal R^2 was used to measure the proportion of outcome variance accounted for by the model (excluding random-effects, i.e., when there is no a-priori knowledge of the expected outcome for each participant). Values vary from 0 to 1, with 1 occurring when the model perfectly predicts the outcome, and 0 occurring when the model only returns the population average.

Finally, a comparison analysis done to compare distress and trauma outcomes in those who completed all 3 surveys against those who dropped out.

Software

All analyses and statistical outputs were produced in the statistical programming language R and the 'tidyverse', 'lme4' and 'ggeffects' packages were used for the mixed-effects models.³⁶⁻³⁸

Patient and Public Involvement

The study team contains frontline doctors from all represented specialties who undertook clinical work throughout the COVID-19 pandemic. This research is in line with recent RCEM research prioritisation and research recommendations.^{39,40}

Role of the funding source

The sponsor and funder had no role at any stage of this work.

Results

Distribution across networks in the UK and Ireland generated 5440 responses. Follow-up responses from the peak and deceleration surveys were 3896 (71·6%) and 3079 (56·6%) respectively (figure 1). The final analysis cohort was 3079 participants, consisting of 1686 (54·8%) from EM, 1114 (36·2%) from Anaesthetics and 526 (17·1%) from ICM, with some participants working across multiple specialities.

The demographic and professional characteristics of the respondent population are summarised in Table 1. The cohort was 51·0% female, with a median age group of 36-40 years, and was representative of all professional grades. Respondents were 63·7% 'White British', 6·2% 'Irish', and '30·1% 'Ethnic Minority'; a full breakdown of ethnicity is provided in the online supplementary hub (<https://github.com/wjchulme/TERN-CERA-study/tree/main/outputs>).^{41,42}

Table 1 Demographic and occupational characteristics of responders who completed all three study phases

	All (N=3079)	Emergency Medicine (N=1686)	Anaesthetics (N=1114)	Intensive Care Medicine (N=526)
Age				
20-25	111 (3·6%)	99 (5·9%)	3 (0·3%)	9 (1·7%)
26-30	737 (24·0%)	471 (28·0%)	184 (16·5%)	130 (24·8%)
31-35	682 (22·2%)	366 (21·7%)	242 (21·8%)	141 (26·9%)
36-40	497 (16·2%)	279 (16·6%)	177 (15·9%)	81 (15·5%)
41-45	406 (13·2%)	220 (13·1%)	156 (14·0%)	55 (10·5%)
46-50	282 (9·2%)	128 (7·6%)	133 (12·0%)	55 (10·5%)
51-55	203 (6·6%)	72 (4·3%)	121 (10·9%)	27 (5·2%)
56-60	107 (3·5%)	34 (2·0%)	63 (5·7%)	19 (3·6%)
>60	49 (1·6%)	14 (0·8%)	33 (3·0%)	7 (1·3%)
Missing	5	3	2	2
Gender				
Male	1455 (48·8%)	774 (47·4%)	542 (50·1%)	272 (53·8%)
Female	1522 (51·0%)	855 (52·4%)	538 (49·7%)	233 (46·0%)
Other	7 (0·2%)	4 (0·2%)	2 (0·2%)	1 (0·2%)
Missing	95	53	32	20
Seniority				
Junior Doctor	1089 (35·4%)	692 (41·0%)	276 (24·8%)	187 (35·6%)
Middle Grade Doctor	660 (21·4%)	357 (21·2%)	230 (20·6%)	129 (24·5%)
Other Senior Doctor	228 (7·4%)	156 (9·3%)	66 (5·9%)	34 (6·5%)
Senior Doctor (Consultant Grade)	1102 (35·8%)	481 (28·5%)	542 (48·7%)	176 (33·5%)
Geographical Region				
East Midlands	177 (5·7%)	78 (4·6%)	84 (7·5%)	24 (4·6%)
East of England	172 (5·6%)	87 (5·2%)	70 (6·3%)	29 (5·5%)
London	454 (14·7%)	319 (18·9%)	103 (9·2%)	42 (8·0%)
North East	132 (4·3%)	68 (4·0%)	47 (4·2%)	30 (5·7%)
North West	334 (10·8%)	149 (8·8%)	141 (12·7%)	78 (14·8%)
South East	355 (11·5%)	229 (13·6%)	105 (9·4%)	48 (9·1%)
South West	430 (14·0%)	208 (12·3%)	167 (15·0%)	76 (14·4%)
West Midlands	183 (5·9%)	89 (5·3%)	78 (7·0%)	44 (8·4%)
Yorkshire and the Humber	212 (6·9%)	90 (5·3%)	102 (9·2%)	55 (10·5%)
Northern Ireland	87 (2·8%)	41 (2·4%)	34 (3·1%)	20 (3·8%)
Scotland	253 (8·2%)	159 (9·4%)	80 (7·2%)	32 (6·1%)

Wales	92 (3·0%)	21 (1·2%)	62 (5·6%)	21 (4·0%)
Dublin	111 (3·6%)	82 (4·9%)	21 (1·9%)	16 (3·0%)
Rest of Ireland	87 (2·8%)	66 (3·9%)	20 (1·8%)	11 (2·1%)
Nation				
England	2449 (79·5%)	1317 (78·1%)	897 (80·5%)	426 (81·0%)
Northern Ireland	87 (2·8%)	41 (2·4%)	34 (3·1%)	20 (3·8%)
Republic of Ireland	198 (6·4%)	148 (8·8%)	41 (3·7%)	27 (5·1%)
Scotland	253 (8·2%)	159 (9·4%)	80 (7·2%)	32 (6·1%)
Wales	92 (3·0%)	21 (1·2%)	62 (5·6%)	21 (4·0%)
Ethnicity				
White British	1888 (63·7%)	949 (58·4%)	755 (70·3%)	338 (67·1%)
Irish	185 (6·2%)	118 (7·3%)	51 (4·7%)	33 (6·5%)
Ethnic minority	893 (30·1%)	557 (34·3%)	268 (25·0%)	133 (26·4%)
Missing	113	62	40	22
Redeployed				
Yes	249 (8·1%)	47 (2·8%)	196 (17·6%)	20 (3·8%)
No	2824 (91·9%)	1636 (97·2%)	916 (82·4%)	504 (96·2%)
Missing	6	3	2	2

Primary Outcomes

General Health Questionnaire-12

The prevalence of psychological distress, as defined by scores >3 on the GHQ-12 0-0-1-1 scoring method, was 44·7% (n=1334) in the acceleration survey, 36·9% (n=1098) at peak and 31·5% (n=918) during the deceleration phase. Median GHQ-12 scores were 13·0 (Q1-Q3, 10·0-17·0), 13·0 (Q1-Q3, 9·0-16·0) and 12·0 (Q1-Q3, 9·0-16·0) respectively (figure 2), and mean scores were 13·7, 13·2 and 12·9 across the acceleration, peak and deceleration surveys. Median distress scores were higher in the Anaesthetic and ICM cohorts at the acceleration phase when compared to EM, but these decreased in all three groups throughout the first pandemic wave.

Impact of Events Scale-Revised

The prevalence of psychological trauma, as defined by a score of >24 on the IES-R, was 23·7% (n=647) at peak and 17·7% (n=484) at deceleration. The prevalence of 'probable PTSD', as defined by a score of >33 was 12·6% (n=343) at peak and 10·1% (n=276) at deceleration. During the peak phase, prevalence of trauma (>24) was 24·9% (n=378) in EM, 21·5% (n=204) in anaesthetics and 24·9% (n=117) in ICM. Prevalence of 'probable PTSD' (>33) was higher in EM (13·9%, n=211) and ICM (13·6%, n= 64) when compared to Anaesthetics (10·8%, n=103). During the deceleration phase, prevalence of trauma (>24) decreased to 19·7% (n=93) in ICM and 18·7% (n=285) in EM. 'Probable PTSD' (>33) decreased to 11·1% (n=169) in EM, compared to 10·8% (n=51) in ICM and 8·8% (n=85) in Anaesthetics. The median IES-R was highest in the peak survey at 13 (Q1-Q3, 5-24), and 9 (Q1-Q3, 2-19) in the deceleration survey (see table 2 and figure 3).

Table 2. GHQ-12 and IES-R Scores for participants who responded to all 3 survey phases

	All (N=3079)	Emergency Medicine (N=1686)	Anaesthetics (N=1114)	Intensive Care Medicine (N=526)
Acceleration				

GHQ-12 (0123 score)				
Mean	13·7	13·3	14·4	14·0
Median (Q1, Q3)	13·0 (10·0, 17·0)	13·0 (10·0, 16·0)	14·0 (11·0, 18·0)	14·0 (10·2, 17·0)
GHQ-12 (0011 > 3)				
> 3	1334 (44·7%)	667 (40·7%)	542 (50·2%)	253 (49·6%)
N-Missing	92	48	34	16
Peak				
GHQ-12 (0123 score)				
Mean	13·2	12·8	13·6	13·6
Median (Q1, Q3)	13·0 (9·0, 16·0)	12·0 (9·0, 16·0)	13·0 (10·0, 17·0)	13·0 (10·0, 17·0)
GHQ-12 (0011 > 3)				
> 3	1098 (36·9%)	543 (33·3%)	454 (42·3%)	211 (41·1%)
N-Missing	105	56	40	13
IES-R score				
Mean	16·3	16·7	15·8	17·2
Median (Q1, Q3)	13·0 (5·0, 24·0)	13·0 (5·0, 24·0)	13·0 (6·0, 23·0)	14·0 (6·0, 24·0)
IES-R > 24				
IES-R-0123 > 24	647 (23·7%)	378 (24·9%)	204 (21·5%)	117 (24·9%)
IES-R > 33				
IES-R-0123 > 33	343 (12·6%)	211 (13·9%)	103 (10·8%)	64 (13·6%)
N-Missing	349	165	163	57
Deceleration				
GHQ-12 (0123 score)				
Mean	12·9	12·8	13·0	13·1
Median (Q1, Q3)	12·0 (9·0, 16·0)	12·0 (9·0, 16·0)	12·0 (9·0, 16·0)	12·0 (9·0, 17·0)
GHQ-12 (0011 > 3)				
> 3	918 (31·5%)	486 (30·2%)	340 (32·6%)	172 (34·6%)
N-Missing	165	78	71	29
IES-R score				
Mean	13·2	13·6	12·6	14·2
Median (Q1, Q3)	9·0 (2·0, 19·0)	9·0 (2·0, 20·0)	8·0 (2·0, 18·0)	9·0 (3·0, 20·0)
IES-R > 24				
IES-R-0123 > 24	484 (17·7%)	285 (18·7%)	159 (16·5%)	93 (19·7%)
IES-R > 33				
IES-R-0123 > 33	276 (10·1%)	169 (11·1%)	85 (8·8%)	51 (10·8%)
N-Missing	344	164	153	53

Secondary Outcomes

Risk Factors for Psychological Distress (GHQ-12) and Trauma (IES-R)

The overall strength of the relationship between participant factors and the two outcome measures, psychological distress and trauma, is summarised using Nagakawa's marginal R^2 (figures 4+5). The form of these univariable relationships is described graphically for the five variables with the highest R^2 values in figures 6 a-f. Graphs for the remaining variables are reported in online supplementary hub (<https://github.com/wjchulme/TERN-CERA-study/tree/main/outputs>).

Personal and Professional variables predicting distress (GHQ-12)

Worry of infecting family members due to clinical work ($R^2 = 0·06$) and worry of personal infection ($R^2 = 0·05$) were the two variables most strongly associated with distress. Figures 6a and 6b report the mean

GHQ-12-score for the levels within this variable. Those that were 'extremely worried' about infecting family had a mean GHQ-12-modelled score of 15.3 (95% CI, 15.0, 15.6), 15.1 (95% CI, 14.8, 15.5) and 14.6 (95% CI, 14.3, 15.0) during the acceleration, peak and deceleration respectively, compared with mean scores of 13.7, 13.2 and 12.9 respectively for all participants. For those who were 'extremely worried' about personal infection, the mean GHQ-12 modelled score was 16.6 (95% CI, 16.1, 17.1) during the acceleration period, compared with 10.9 (95% CI, 9.7, 12.1) for those who were 'not worried at all' about being infected. For the mean GHQ-12 modelled score for each of the other variables see the online link for the figures and values (<https://github.com/wjchulme/TERN-CERA-study/tree/main/outputs>).

Personal and Professional variables predicting trauma (IES-R)

For trauma, worry of infection of family members due to clinical role had the highest R^2 value ($R^2=0.10$). Mean IES-R modelled score for those who were 'extremely worried' about infecting family was 23.0 (95% CI, 22.2, 23.8) during the peak compared to 10.0 (95% CI, 7.8, 12.2) for those who were 'not worried at all' during the peak (Fig 6c). This is significantly higher than the reported mean IES-R overall of 16.3.

Concern that COVID-19 would exacerbate symptoms of an established mental health condition ($R^2 = 0.06$) had the second highest R^2 value. Peak IES-R mean modelled scores were 23.3 (95% CI, 22.1, 24.4) in those who agreed with this statement compared to 15.2 (95% CI, 14.7, 15.7) in those who disagreed. Deceleration mean IES-R modelled scores remained high for those who agreed, 22.3 (95% CI, 21.1, 23.6). (Figure 6d)

Worry relating to personal infection due to clinical role ($R^2 = 0.06$) was again strongly associated with trauma. Figure 6e displays the mean IES-R modelled scores and demonstrates the peak (24.0 (95% CI, 22.5, 25.4)) and deceleration (20.3 (95% CI, 18.7, 21.8)) outcomes in participants who were 'extremely worried' compared to those who were 'not worried at all' during the peak (11.3 (95% CI 8.6, 14.0)) and deceleration (10.0 (95% CI 8.0, 12.0)).

Whilst ethnicity was not strongly associated with distress, it was a stronger predictor of trauma ($R^2 = 0.03$). Mean modelled trauma scores for 'Ethnic Minority' participants at peak was 18.8 (95% CI, 17.8, 19.8), compared to 'White British' participants of 15.1 (95% CI, 14.5, 15.8). (Figure 6f) For the mean IES-R modelled scores for each of the other variables see online link for the figures and values. (<https://github.com/wjchulme/TERN-CERA-study/tree/main/outputs>)

Incidence of self-reported COVID-19 infection and isolation

By the deceleration phase of the pandemic 6.9% (n=212) of respondents had received a positive diagnosis of COVID-19 and 0.4% (n=12) had been admitted to hospital. A positive diagnosis did not have a significant effect in prediction of trauma ($R^2=0.014$).

Regional and national variation of psychological distress and trauma

The region in which participants worked was more valuable for predicting trauma ($R^2 = 0.034$), than for distress ($R^2 = 0.016$). The mean modelled score of the different regions within the UK and Republic of Ireland on IES-R is demonstrated in figure 7.

Drop-out by GHQ-12 and IES-R

Response rate for the peak and deceleration surveys was 71.6% and 56.6% respectively. There was no significant difference in either the GHQ-12 or IES-R scores between those who dropped out and those who remained in the study (see online supplement).

Discussion

In this prospective longitudinal survey of 3079 frontline doctors, the prevalence of psychological distress reached 44.7% during the acceleration phase, and reached 23.7% for trauma during the peak phase - these figures were substantially higher than for the general population.⁴³ For psychological distress, rates declined through peak and deceleration phases of the first wave to a level comparable to pre-pandemic levels.⁴⁴ Prevalence of 'probable PTSD' was 12.6% at peak and 10.1% at deceleration, demonstrating a degree of natural recovery.^{45,46} However, just less than a quarter experienced sub-threshold post-trauma symptoms 30 days following the pandemic peak.

Personal factors were the most powerful predictors of both psychological distress and trauma. The most significant predictors relate to familial safety, personal safety, and established mental health conditions. These findings support aggregated data in recent reviews and meta-analyses on the key predictors of psychological distress in disaster or infectious outbreak settings.^{1,7,21-23} However, it cannot be ignored that the psychological harm associated with both familial and personal safety may potentially be explained by the perceived (and reported) inadequate provision of PPE to frontline workers.^{47,48} This is an area where improvements must be made in order to mitigate against future physical and psychological harms that novel pathogens present.

While most findings are consistent with existing research, our study also identifies ethnicity as a novel, key predictor of trauma.⁴⁹⁻⁵¹ This is unsurprising given higher rates of reported mortality in ethnic minority groups with this particular pandemic.⁵² However the nature and direction of relationship between these risk factors and poorer outcomes is undoubtedly complex. Ongoing work continues to seek further understanding in this area.⁵³

Rates of trauma were high across all three specialty groups. One in four doctors met the clinical threshold, with the highest rates seen in EM and ICM. This is likely explained by their clinical roles during the pandemic, in which they were exposed to a higher volume of COVID-19 positive patients compared to Anaesthetic colleagues. However, it is important to note that the rate of trauma seen in Anaesthetics was also of concern. At the deceleration phase, EM doctors had higher rates of 'probable PTSD' (IES-R >33), whereas ICM doctors had a higher prevalence of trauma (IES-R >24). This may

reflect the later peak in ICUs when compared to EM ⁵⁴ or the potential impact of downstream mortality. Further work should explore longer term outcomes in all cohorts.

It is evident from our longitudinal data that vulnerability to poorer psychological outcomes may be predicted by certain characteristics and therefore potentially mitigated through targeted intervention. Studies examining psychiatric outcomes in SARS reflect that psychological distress is likely to persist. Identification of those likely to experience adversity, and interventions to mitigate these, must begin now. ^{8,10,55,56} Without appropriate support and intervention doctors are likely to experience long-term effects on mental health, resulting in increased sickness rates, absenteeism, impaired performance at work, and the development of physical health problems. ^{8,10,12,57,58} Therefore the early identification of ongoing psychological distress will be pivotal in influencing the longer-term mental health of frontline workers. Based on research from COVID-19 and other pandemics, we can be certain that rates and severity of distress will rise following this second wave of the pandemic. We now know that doctors are working on the frontline while carrying the heavy burden of fear of infecting themselves, or critically, family members, while some continue to battle high levels of psychological distress. This distress was evident in the lead up to the first peak, but sustained well beyond this time point. Doctors are continuing to work in very highly pressured, high risk environments with a significant proportion doing so despite clinical levels of distress. Policymakers and professional bodies should urgently seek to develop an overarching 'best practice' pathway to support all healthcare staff in these environments.

While various interventions are recommended specifically for frontline workers there is common agreement in the necessity for basic psychosocial interventions (i.e. sleep hygiene, exercise, health behaviour) to facilitate return to equilibrium ⁵⁹⁻⁶², yet these measures are not always sufficient to ameliorate persistent distress. It is crucial that an overarching 'best practice' pathway and package of care is implemented to help support staff now and for the future. This must be evidence-based, multilevel, starting with the 'individual' level and moving through to 'organisational' level intervention, including (a) mobilisation of formal peer & organisational support structures, (b) mechanisms for recognising and monitoring distress, and (c) offer clear referral pathways to evidence-based interventions. Access to appropriate psychological support is imperative; cognitive behavioural therapy is recommended by the National Institute for health and Care Excellence (NICE) to ameliorate anxiety, depression and PTSD ^{63,64} however further work is needed to ensure these interventions are suitably tailored to the practicalities of shift work and the unique experiences faced by frontline clinicians. With this, there is a responsibility to ensure equality in the provision of care and pathways to access, for this is likely to be necessary for many.

Strengths and Weaknesses

This is a large-scale longitudinal study examining prevalence of psychological distress in doctors in the UK and Ireland, offering a robust and reliable measure of the impact of COVID-19 on the mental health of frontline doctors, and allows comparison with other pandemic mental health trajectories. Due to the three-phase prospective design and extent of data collected, findings from this study can be reliably

used to inform the development of preparations and interventions to mitigate the impact of COVID-19 and future infectious disease outbreaks on mental health in frontline doctors.

However, there are limitations that may influence our findings. The reported rates of distress and trauma do not take account of any pre-existing psychiatric morbidity or historical factors that may predispose doctors to developing mental health difficulties in these circumstances.^{40,65–67} Data was gathered with regards to historical trauma, one of the most significant predictors of mental health difficulties long-term. Furthermore, whilst the sample size is large, any self-reporting measure is open to selection bias. This may have resulted in a biased sample with particularly high or low levels of distress and trauma. However, in the follow-up surveys (peak and deceleration) there was no difference in acceleration distress or trauma scores between those who dropped out and those who continued; yet we are unable to comment on those who declined to participate. Whilst the two primary outcome measures, GHQ-12 and IES-R, have good psychometric properties, there is a concern that survey data may overstate the prevalence of cases when compared to formal diagnostic interviews such as the SCID (The Structured Clinical **Interview** for DSM-IV Axis I Disorders); this is difficult to implement in such large samples, thus we cautiously avoid inference of definite diagnosis.

While the protocol was closely adhered to, variation in regional peaks may have influenced accurate capturing of psychological distress and trauma rates. It is noted that whilst the acceleration phase is study 'baseline', as the pandemic was present and proliferating in the UK at the acceleration phase, it more accurately represents the initial stress associated with a rapidly spreading highly infectious virus of unknown pathogenic origins and no effective treatment; a reasonable response to the context. Future research should continue to follow frontline doctors through the pandemic and beyond, to assess whether the mental health trajectories are similar to other infectious disease pandemics.

Conclusion

Our findings reflect a pattern of elevated distress during the acceleration and peak phase of the current pandemic, some degree of natural recovery and a significant minority continuing to experience residual ongoing distress. It is essential that policymakers and professional bodies seek to prevent future adverse effects through provision of vital equipment to mitigate both physical and psychological harm and the development of clear pathways to effective psychological care. Moving forward, it is essential the COVID-19 pandemic serves as a foundation for significant development and growth in all of these areas and that there is ongoing assessment of the psychological health of healthcare workers both during the pandemic and beyond.

References

- 1 Brooks SK, Dunn R, Sage CAM, Amlôt R, Greenberg N, Rubin GJ. Risk and resilience factors affecting the psychological wellbeing of individuals deployed in humanitarian relief roles after a disaster. *J. Ment. Heal.* 2015. DOI:10.3109/09638237.2015.1057334.
- 2 Liu Q, Luo D, Haase JE, *et al.* The experiences of health-care providers during the COVID-19

- crisis in China: a qualitative study. *Lancet Glob Heal* 2020. DOI:10.1016/S2214-109X(20)30204-7.
- 3 McCabe R, Schmit N, Christen P, *et al.* Adapting hospital capacity to meet changing demands during the COVID-19 pandemic. *BMC Med* 2020. DOI:10.1186/s12916-020-01781-w.
 - 4 Phua J, Weng L, Ling L, *et al.* Intensive care management of coronavirus disease 2019 (COVID-19): challenges and recommendations. *Lancet Respir. Med.* 2020. DOI:10.1016/S2213-2600(20)30161-2.
 - 5 Kursumovic E, Lennane S, Cook TM. Deaths in healthcare workers due to COVID-19: the need for robust data and analysis. *Anaesthesia* 2020; **75**: 989–92.
 - 6 Maunder RG, Lancee WJ, Rourke S, *et al.* Factors associated with the psychological impact of severe acute respiratory syndrome on nurses and other hospital workers in Toronto. *Psychosom. Med.* 2004. DOI:10.1097/01.psy.0000145673.84698.18.
 - 7 Kisely S, Warren N, McMahon L, Dalais C, Henry I, Siskind D. Occurrence, prevention, and management of the psychological effects of emerging virus outbreaks on healthcare workers: rapid review and meta-analysis. *BMJ* 2020. DOI:10.1136/bmj.m1642.
 - 8 Allan SM, Bealey R, Birch J, *et al.* The prevalence of common and stress-related mental health disorders in healthcare workers based in pandemic-affected hospitals: a rapid systematic review and meta-analysis. 2020 DOI:10.1101/2020.05.04.20089862.
 - 9 Roberts T, Daniels J, Hulme W, *et al.* Psychological distress during the acceleration phase of the COVID-19 pandemic: A survey of doctors practising in emergency medicine, anaesthesia and intensive care medicine in the UK and Ireland. *Emerg Med J* 2021; **0**: 1–10.
 - 10 Maunder RG, Lancee WJ, Balderson KE, *et al.* Long-term psychological and occupational effects of providing hospital healthcare during SARS outbreak. *Emerg Infect Dis* 2006. DOI:10.3201/eid1212.060584.
 - 11 Fiksenbaum L, Marjanovic Z, Greenglass ER, Coffey S. Emotional exhaustion and state anger in nurses who worked during the sars outbreak: The role of perceived threat and organizational support. *Can J Community Ment Heal* 2006. DOI:10.7870/cjcmh-2006-0015.
 - 12 Arora M, Asha S, Chinnappa J, Diwan AD. Review article: Burnout in emergency medicine physicians. *EMA - Emerg. Med. Australas.* 2013; **25**: 491–5.
 - 13 Lai J, Ma S, Wang Y, *et al.* Factors Associated With Mental Health Outcomes Among Health Care Workers Exposed to Coronavirus Disease 2019. *JAMA Netw open* 2020; **3**: e203976.
 - 14 Di Tella M, Romeo A, Benfante A, Castelli L. Mental health of healthcare workers during the <scp>COVID</scp> -19 pandemic in Italy. *J Eval Clin Pract* 2020; **26**: 1583–7.
 - 15 Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsis E, Katsaounou P. Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain Behav Immun* 2020. DOI:10.1016/j.bbi.2020.05.026.
 - 16 Benfante A, Di Tella M, Romeo A, Castelli L. Traumatic Stress in Healthcare Workers During COVID-19 Pandemic: A Review of the Immediate Impact. *Front Psychol* 2020; **11**: 569935.
 - 17 Clap for Carers: UK in “emotional” tribute to NHS and care workers - BBC News.

- <https://www.bbc.co.uk/news/uk-52058013> (accessed Nov 20, 2020).
- 18 The Lancet. COVID-19: protecting health-care workers. *Lancet*. 2020. DOI:10.1016/S0140-6736(20)30644-9.
 - 19 Coronavirus: Mental health of NHS staff at long-term risk - BBC News.
<https://www.bbc.co.uk/news/health-52528619> (accessed Nov 20, 2020).
 - 20 Halpern J, Maunder RG, Schwartz B, Gurevich M. Identifying risk of emotional sequelae after critical incidents. *Emerg Med J* 2011. DOI:10.1136/emj.2009.082982.
 - 21 Brooks SK, Dunn R, Amlôt R, Rubin GJ, Greenberg N. A Systematic, Thematic Review of Social and Occupational Factors Associated with Psychological Outcomes in Healthcare Employees during an Infectious Disease Outbreak. *J Occup Environ Med* 2018. DOI:10.1097/JOM.0000000000001235.
 - 22 Lancee WJ, Maunder RG, Goldbloom DS. Prevalence of psychiatric disorders among Toronto hospital workers one to two years after the SARS outbreak. *Psychiatr Serv* 2008. DOI:10.1176/ps.2008.59.1.91.
 - 23 Ozer EJ, Best SR, Lipsey TL, Weiss DS. Predictors of posttraumatic stress disorder and symptoms in adults: A meta-analysis. *Psychol. Bull.* 2003; **129**: 52–73.
 - 24 Roberts T, Daniels J, Hulme W, *et al.* COVID-19 emergency response assessment study: a prospective longitudinal survey of frontline doctors in the UK and Ireland: study protocol. *BMJ Open* 2020. DOI:10.1136/bmjopen-2020-039851.
 - 25 Eysenbach G. Improving the Quality of Web Surveys: The Checklist for Reporting Results of Internet E-Surveys (CHERRIES). *J Med Internet Res* 2004; **6**: e34.
 - 26 Harris PA, Taylor R, Minor BL, *et al.* The REDCap consortium: Building an international community of software platform partners. *J Biomed Inform* 2019; **95**: 103208.
 - 27 Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009; **42**: 377–81.
 - 28 Holloway R, Rasmussen SA, Zaza S, Cox N, Jernigan D. Updated Preparedness and Response Framework for Influenza Pandemics. 2014.
<https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6306a1.htm> (accessed April 8, 2020).
 - 29 Goldberg D, Williams P. A user's guide to the General Health Questionnaire. London: GL Assessment, 1988.
 - 30 Goldberg DP, Gater R, Sartorius N, *et al.* The validity of two versions of the GHQ in the WHO study of mental illness in general health care. *Psychol Med* 1997. DOI:10.1017/S0033291796004242.
 - 31 Goldberg DP, Oldehinkel T, Ormel J. Why GHQ threshold varies from one place to another. *Psychol Med* 1998. DOI:10.1017/S0033291798006874.
 - 32 Christianson S, Marren J. The Impact of Event Scale - Revised (IES-R). *Medsurg Nurs* 2012.
 - 33 Asukai N, Kato H, Kawamura N, *et al.* Reliability and validity of the Japanese-language version of the Impact of Event Scale-Revised (IES-R-J): Four studies of different traumatic events. *J Nerv Ment Dis* 2002. DOI:10.1097/00005053-200203000-00006.

- 34 Creamer M, Bell R, Failla S. Psychometric properties of the Impact of Event Scale - Revised. *Behav Res Ther* 2003. DOI:10.1016/j.brat.2003.07.010.
- 35 Nakagawa S, Johnson PCD, Schielzeth H. The coefficient of determination R² and intra-class correlation coefficient from generalized linear mixed-effects models revisited and expanded. *J R Soc Interface* 2017. DOI:10.1098/rsif.2017.0213.
- 36 Wickham H, Averick M, Bryan J, *et al*. Welcome to the Tidyverse. *J Open Source Softw* 2019; **4**: 1686.
- 37 Bates D, Mächler M, Bolker BM, Walker SC. Fitting linear mixed-effects models using lme4. *J Stat Softw* 2015. DOI:10.18637/jss.v067.i01.
- 38 Lüdtke D. ggeffects: Tidy Data Frames of Marginal Effects from Regression Models. *J Open Source Softw* 2018; **3**: 772.
- 39 Smith J, Keating L, Flowerdew L, *et al*. An Emergency Medicine Research Priority Setting Partnership to establish the top 10 research priorities in emergency medicine. *Emerg Med J* 2017. DOI:10.1136/emmermed-2017-206702.
- 40 Cottey L, Roberts T, Graham B, *et al*. Need for recovery amongst emergency physicians in the UK and Ireland: a cross-sectional survey. *BMJ Open* 2020.
- 41 Khunti K, Routen A, Pareek M, Treweek S, Platt L. The language of ethnicity. *BMJ* 2020; **371**: m4493.
- 42 Bunglawala Z (Race DU, Office) C. Please, don't call me BAME or BME! 2019. <https://civilservice.blog.gov.uk/2019/07/08/please-dont-call-me-bame-or-bme/> (accessed Oct 12, 2020).
- 43 Rettie H, Daniels J. Coping and Tolerance of Uncertainty: Predictors and Mediators of Mental Health During the COVID-19 Pandemic. *Am Psychol* 2020. DOI:10.1037/amp0000710.
- 44 Kinman G, Teoh K. What could make a difference to the mental health of UK doctors? A review of the research evidence. 2018.
- 45 Morina N, Wicherts JM, Lobbrecht J, Priebe S. Remission from post-traumatic stress disorder in adults: A systematic review and meta-analysis of long term outcome studies. *Clin. Psychol. Rev.* 2014. DOI:10.1016/j.cpr.2014.03.002.
- 46 Kessler RC, Sonnega A, Bromet E, Hughes M, Nelson CB. Posttraumatic Stress Disorder in the National Comorbidity Survey. *Arch Gen Psychiatry* 1995. DOI:10.1001/archpsyc.1995.03950240066012.
- 47 McKee M. England's PPE procurement failures must never happen again. *BMJ*. 2020. DOI:10.1136/bmj.m2858.
- 48 Godlee F. Covid-19: weathering the storm. *BMJ* 2020; **368**: m1199.
- 49 Chew NWS, Lee GKH, Tan BYQ, *et al*. A multinational, multicentre study on the psychological outcomes and associated physical symptoms amongst healthcare workers during COVID-19 outbreak. *Brain Behav Immun* 2020. DOI:10.1016/j.bbi.2020.04.049.
- 50 Berger W, Coutinho ESF, Figueira I, *et al*. Rescuers at risk: A systematic review and meta-regression analysis of the worldwide current prevalence and correlates of PTSD in rescue workers. *Soc. Psychiatry Psychiatr. Epidemiol.* 2012. DOI:10.1007/s00127-011-0408-2.

- 51 Perrin MA, DiGande L, Wheeler K, Thorpe L, Farfel M, Brackbill R. Differences in PTSD prevalence and associated risk factors among World Trade Center disaster rescue and recovery workers. *Am J Psychiatry* 2007. DOI:10.1176/appi.ajp.2007.06101645.
- 52 Coronavirus (COVID-19) related deaths by ethnic group, England and Wales - Office for National Statistics.
<https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/coronavirusrelateddeathsbyethnicgroupenglandandwales/2march2020to10april2020> (accessed Nov 20, 2020).
- 53 Iacobucci G. Covid-19: Increased risk among ethnic minorities is largely due to poverty and social disparities, review finds. *BMJ* 2020; **371**. DOI:10.1136/bmj.m4099.
- 54 Doidge JC, Mouncey PR, Thomas K, *et al*. Trends in Intensive Care for Patients with COVID-19 in England, Wales and Northern Ireland. 2020; published online Aug 11.
DOI:10.20944/preprints202008.0267.v1.
- 55 Mak IWC, Chu CM, Pan PC, Yiu MGC, Chan VL. Long-term psychiatric morbidities among SARS survivors. *Gen Hosp Psychiatry* 2009. DOI:10.1016/j.genhosppsych.2009.03.001.
- 56 Lee AM, Wong JGWS, McAlonan GM, *et al*. Stress and psychological distress among SARS survivors 1 year after the outbreak. *Can J Psychiatry* 2007.
DOI:10.1177/070674370705200405.
- 57 McAlonan GM, Lee AM, Cheung V, *et al*. Immediate and sustained psychological impact of an emerging infectious disease outbreak on health care workers. *Can J Psychiatry* 2007.
DOI:10.1177/070674370705200406.
- 58 Stuijzand S, Deforges C, Sandoz V, *et al*. Psychological impact of an epidemic/pandemic on the mental health of healthcare professionals: A rapid review. *BMC Public Health* 2020; **20**: 1230.
- 59 Greenberg N, Docherty M, Gnanapragasam S, Wessely S. Managing mental health challenges faced by healthcare workers during covid-19 pandemic. *BMJ*. 2020.
DOI:10.1136/bmj.m1211.
- 60 Que J, Shi L, Deng J, *et al*. Psychological impact of the covid-19 pandemic on healthcare workers: A cross-sectional study in China. *Gen Psychiatry* 2020. DOI:10.1136/gpsych-2020-100259.
- 61 Maben J, Bridges J. Covid-19: Supporting nurses' psychological and mental health. *J. Clin. Nurs*. 2020. DOI:10.1111/jocn.15307.
- 62 Heath C, Sommerfield A, von Ungern-Sternberg BS. Resilience strategies to manage psychological distress among healthcare workers during the COVID-19 pandemic: a narrative review. *Anaesthesia*. 2020. DOI:10.1111/anae.15180.
- 63 Post-traumatic stress disorder NICE guideline. 2018 www.nice.org.uk/guidance/ng116 (accessed Nov 23, 2020).
- 64 Kendrick T, Pilling S. Common mental health disorders - Identification and pathways to care: NICE clinical guideline. *Br. J. Gen. Pract.* 2012; **62**: 47–9.
- 65 Graham B, Cottey L, Smith J, Mills M, Latour J. Measuring 'Need for Recovery' as an indicator

- of staff wellbeing in the Emergency Department—a survey study. *Emerg Med J* 2020.
- 66 Basu S, Qayyum H, Mason S. Occupational stress in the ED: A systematic literature review. *Emerg Med J* 2017; **34**: 441–7.
- 67 Schneider A, Weigl M. Associations between psychosocial work factors and provider mental well-being in emergency departments: A systematic review. *PLoS One*. 2018; **13**. DOI:10.1371/journal.pone.0197375.

Acknowledgements

The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, the Department of Health or the Royal Colleges involved in survey distribution.

The authors would like to acknowledge Mai Baquedano, at the University of Bristol, for her support with REDCap, GL Assessments for providing the licence for the GHQ-12 free of charge and Simon O'Hare, Data and Insight Manager, General Medical Council

Author Contributions

The corresponding author attests that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted. Tom Roberts (TR) conceived the idea for the study. TR, Edd Carlton (EC), Jo Daniels (JD), Mark Lyttle (ML), and Blair Graham (BG) were responsible for the initial study design, which was refined with the help of Katie Samuel (KS), Charles Reynard (CR), Robert Hirst (RH), Michael Barrett (MB) and William Hulme (WH). Expert advice on psychological assessment scores was provided by JD. WH provided the statistical plan. TR lead the dissemination of the study in UK Adult Emergency Departments (ED), ML lead the dissemination of the study in UK and Ireland Paediatric EDs, KS lead the dissemination of the study in UK Anaesthetic and ICU Departments, MB lead the dissemination of the study in Ireland EDs, along with John Cronin, James Foley and Etimbuk Umana. Joao Vinagre lead the dissemination in Ireland ICUs and Anaesthetic Departments. TR coordinated study set-up, finalisation of the study surveys and finalisations of study protocols. All authors contributed to the final study design and protocol development, critically revised successive drafts of the manuscript and approved the final version. The study management group is responsible for the conduct of the study.

Funding

The Chief Investigator is directly funded as a research fellow by the Royal College of Emergency Medicine. The GHQ-12 is being used under licence from GL assessments; the fee for use of this instrument within all three surveys has been waived. Dr Carlton is a National Institute for Health Research Advanced Fellow. The study has direct funding from RCEM. Grant code: G/2020/1.

Competing interests

Many of the authors have been working as frontline clinicians during the COVID-19 pandemic. They have no competing interests to declare.

Data Sharing

Deidentified participant data will be made available for 2 years post publication. Requests for access will require HRA and ethical approval and decisions regarding data sharing will be made after discussion with the study senior authors. Statistical code and study figures are available directly from: <https://github.com/wjchulme/TERN-CERA-study/tree/main/outputs>.

Trainee Emergency Research Network (TERN) Collaborators:

L	Kane	Aberdeen Royal Infirmary	R	Hannah	Royal Alexandra Children's Hospital
L	Mackenzie	Addenbrooke's Hospital, Cambridge	A	Corfield	Royal Alexandra Hospital, Scotland
S	Sharma Hajela	Addenbrooke's Hospital, Cambridge	J	Maney	Royal Belfast Hospital for Sick Children
J	Phizacklea	Addenbrooke's Hospital, Cambridge	D	Metcalfe	Royal Berkshire Hospital
K	Malik	Addenbrooke's Hospital, Cambridge	S	Timmis	Royal Berkshire Hospital
N	Mathai	Aintree University Hospital	C	Williams	Royal Bolton Hospital
A	Sattout	Aintree University Hospital	R	Newport	Royal Bolton Hospital
S	Messahel	Alder Hey Children's Hospital, Liverpool	D	Bawden	Royal Cornwall Hospital
E	Fadden	Alder Hey Children's Hospital, Liverpool	A	Tabner	Royal Derby Hospital
R	McQuillan	Altnagelvin Area Hospital, N. Ireland	H	Malik	Royal Devon and Exeter Hospital
B	O'Hare	Antrim Area Hospital, N. Ireland	C	Roe	Royal Devon and Exeter Hospital
P	Turton	Arrowe Park Hospital, Merseyside	D	McConnell	Royal Devon and Exeter Hospital
S	Lewis	Arrowe Park Hospital, Merseyside	F	Taylor	Royal Free London
D	Bewick	Barnsley Hospital	R	Ellis	Royal Glamorgam Hospital, Wales
R	Taylor	Bath Royal United Hospital	S	Morgan	Royal Gwent Hospital, Wales
I	Hancock	Bath Royal United Hospital	L	Barnicott	Royal Hampshire County Hospital
D	Manthalapo Ramesh Babu	Bedford Hospital, Bedfordshire	S	Foster	Royal Hospital for Children, Glasgow
S	Hartshorn	Birmingham Children's Hospital	J	Browning	Royal Hospital for Sick Children Royal Hospital for Sick Children, Edinburgh
M	Williams	Birmingham Children's Hospital	L	McCrae	
A	Charlton	Bradford Royal Infirmary	E	Godden	Royal Infirmary Hospital, Edinburgh
L	Somerset	Bristol Royal Hospital for Children	A	Saunders Lawrence- Ball	Royal Infirmary Hospital, Edinburgh
C	Munday	Bristol Royal Hospital for Children	A		Royal Liverpool University Hospital
A	Turner	Bristol Royal Hospital for Children	R	House	Royal Liverpool University Hospital
R	Sainsbury	Bristol Royal Infirmary	J	Muller	Royal London Hospital
E	Williams	Bristol Royal Infirmary	I	Skene	Royal London Hospital
S	Patil	Chelsea & Westminster Hospital	M	Lim	Royal London Hospital
R	Stewart	Chelsea & Westminster Hospital	H	Millar	Royal Manchester Children's Hospital
M	Winstanley	Chelsea & Westminster Hospital	A	Rai	Royal Manchester Children's Hospital
N	Tambe	Chesterfield Royal hospital	K	Challen	Royal Preston Hospital
C	Magee	City Hospital, Birmingham	S	Currie	Royal Preston Hospital
D	Raffo	Craigavon Area Hospital, N. Ireland	M	Elkanzi	Royal Stoke University Hospital
D	Mawhinney	Craigavon Area Hospital, N. Ireland	T	Perry	Royal Surrey County Hospital
B	Taylor	Cumberland Infirmary, Cumbria	W	Kan	Royal Surrey County Hospital
T	Hussan	Darlington Memorial Hospital	L	Brown	Royal Sussex County Hospital
G	Pells	Darlington Memorial Hospital	M	Cheema	Royal Sussex County Hospital
F	Barham	Derriford Hospital, Plymouth	A	Clarey	Royal Victoria Hospital
F	Wood	Derriford Hospital, Plymouth	A	Gulati	Royal Victoria Infirmary
C	Szekeres	East Surrey Hospital	K	Webster	Royal Victoria Infirmary
R	Greenhalgh	East Surrey Hospital	A	Howson	Salford Royal NHS Foundation Trust

S	Marimuthu	Eastbourne Hospital	R	Doonan	Salford Royal NHS Foundation Trust
R	Macfarlane	Epsom and St Helier Hospitals	C	Magee	Sandwell Hospital
M	Alex	Evelina Children's Hospital, London	A	Trimble	Sheffield Children's Hospital
B	Shrestha	Frimley Park Hospital	C	O'Connell	Sheffield Children's Hospital
L	Stanley	Gloucester Royal Hospital	R	Wright	Southampton General
J	Gumley	Gloucester Royal Hospital	E	Colley	Southmead Hospital, Bristol
K	Thomas	Gloucester Royal Hospital	C	Rimmer	Southport Hospital, Merseyside
M	Anderson	Great North Children's, Newcastle	S	Pintus	Southport Hospital, Merseyside
C	Weegenaar	Great Western Hospital, Swindon	H	Jarman	St George's Hospital, London
J	Lockwood	Harrogate Hospital	V	Worsnop	St George's Hospital, London
T	Mohamed	Heartlands's Hospital, Birmingham	S	Collins	St Helier Hospital
S	Ramraj	Hillingdon Hospital, London	M	Colmar	St John's Hospital, Livingston
M	Mackenzie	Homerton Hospital, London	N	Masood	St John's Hospital, Livingston
A	Robertson	Homerton Hospital, London	R	McLatchie	St John's Hospital, Livingston
W	Niven	Homerton Hospital, London	A	Peasley	Stepping Hill Hospital
M	Patel	Homerton Hospital, London	S	Rahman	Stoke Mandeville Hospital
S	Subramaniam	Horton General Hospital, Banbury	N	Mullen	South Tyneside and Sunderland NHS Trust
C	Holmes	Huddersfield Royal Infirmary	L	Armstrong	The Royal Berkshire Hospital, Reading
S	Bongale	Inverclyde Royal Hospital	A	Hay	The Whittington Hospital, London
U	Bait	Ipswich Hospital	R	Mills	The Whittington Hospital, London
S	Nagendran	Ipswich Hospital	J	Lowe	Torbay Hospital, Devon
S	Rao	Ipswich Hospital	H	Raybould	Torbay Hospital, Devon
F	Mendes	James Paget Hospital	A	Ali	Torbay Hospital, Devon
P	Singh	John Radcliffe Hospital, Oxford	P	Cuthbert	Ulster Hospital Dundonald, N. Ireland
S	Subramaniam	John Radcliffe Hospital, Oxford	S	Taylor	University College London Hospital
T	Baron	John Radcliffe Hospital, Oxford	V	Talwar	University College London Hospital
C	Ponmani	King George Hospital	Z	Al-Janabi	University Hospital Ayr, Scotland
M	Depante	King's College Hospital, London	C	Leech	University Hospital Coventry
R	Sneep	King's College Hospital, London	J	Turner	University Hospital Coventry
A	Brookes	King's College Hospital, London	L	McKechnie	University Hospital Crosshouse, Scotland
S	Williams	King's College Hospital, London	B	Mallon	University Hospital Crosshouse, Scotland
A	Rainey	King's College Hospital, London	J	McLaren	University Hospital Crosshouse, Scotland
J	Brown	Kingston Hospital, London	Y	Moulds	University Hospital Crosshouse, Scotland
N	Marriage	Kingston Hospital, London	L	Dunlop	University Hospital Hairmyres, Scotland
S	Manou	Leeds General Infirmary	FM	Burton	University Hospital Hairmyres, Scotland
S	Hart	Leeds General Infirmary	S	Keers	University Hospital Lewisham, London
M	Elsheikh	Leeds General Infirmary	L	Robertson	University Hospital Lewisham, London
L	Cocker	Leicester Royal Infirmary	D	Craver	University Hospital Lewisham, London
MH	Elwan	Leicester Royal Infirmary	N	Moultrie	University Hospital Monklands, Scotland
K L	Vincent	Leicester Royal Infirmary	O	Williams	University Hospital of North Tees
C	Nunn	Leicester Royal Infirmary	S	Purvis	University Hospital of North Tees
N	Sarja	Lister Hospital, Stevenage	M	Clark	University Hospital of North Tees
M	Viegas	Luton & Dunstable Hospital	C	Davies	University Hospital of Wales, Cardiff
E	Wooffinden	Manchester Royal Infirmary	S	Foreman	University Hospital of Wales, Cardiff
C	Reynard	Manchester Royal Infirmary	C	Ngua	University Hospital of Wales, Cardiff
N	Cherian	Manchester Royal Infirmary	D	George	University Hospital of Wales, Cardiff

A	Da-Costa	Medway NHS Foundation Trust	J	Morgan	University Hospital of Wales, Cardiff
S	Duckitt	Medway NHS Foundation Trust	D	George	University Hospital of Wales, Cardiff
J	Bailey	Milton Keynes University Hospital	N	Hoskins	University Hospital of Wales, Cardiff
L	How	Milton Keynes University Hospital	J	Fryer	University Hospital Southampton
T	Hine	Milton Keynes University Hospital	R	Wright	University Hospital Southampton
F	Ihsan	Milton Keynes University Hospital	L	Frost	University Hospital Southampton
H	Abdullah	Milton Keynes University Hospital	P	Ellis	University Hospital Southampton
K	Bader	Milton Keynes University Hospital	A	Mackay	University Hospital Wishaw, Scotland
S	Pradhan	Milton Keynes University Hospital	K	Gray	Victoria Hospital, Kirkcaldy, Scotland
M	Manoharan	Milton Keynes University Hospital	M	Jacobs	Watford General Hospital
C	Battle	Morrison Hospital, Wales	I	Muslim Veetil Asif	West Middlesex university hospital
L	Kehler	Wolverhampton NHS Trust	P	Amiri	West Middlesex university hospital
R	Muswell	Newham University Hospital, London	S	Shrivastava	West Middlesex university hospital
M	Bonsano	Newham University Hospital, London	F	Raza	West Middlesex university hospital
J	Evans	Norfolk and Norwich Hospitals	S	Wilson	Wexham Park Hospital
E	Christmas	North Hampshire Hospital, Basingstoke	M	Riyat	Wexham Park Hospital
K	Knight	North Middlesex Hospital, London	H	Knott	Wexham Park Hospital
L	O'Rourke	North Tees Hospital, Stockton on Tees	M	Ramazany	Whiston Hospital, Merseyside
K	Adeboye	North Tees Hospital, Stockton on Tees	S	Langston	Whiston Hospital, Merseyside
K	Iftikhar	Northern General Hospital, Sheffield	N	Abela	Whiston Hospital, Merseyside
R	Evans	Northern General Hospital, Sheffield	L	Robinson	Whittington Hospital, London
R	Darke	Northumbria Specialist Emergency Hospital	D	Maasdorp	Whittington Hospital, London
R	Freeman	Northumbria Specialist Emergency Hospital	H	Murphy	Whittington Hospital, London
E	Grocholski	Northwick Park Hospital, London	H	Edmundson	Whittington Hospital, London
K	Kaur	Peterborough City Hospital	R	Das	Whittington Hospital, London
H	Cooper	Peterborough City Hospital	C	Orjioke	Whittington Hospital, London
M	Mohammad	Princess Royal Hospital, London	D	Worley	Whittington Hospital, London
L	Harwood	Princess Royal Hospital, London	W	Collier	Whittington Hospital, London
K	Lines	Queen Alexandra Hospital, Portsmouth	J	Everson	Whittington Hospital, London
C	Thomas	Queen Alexandra Hospital, Portsmouth	N	Maleki	Whittington Hospital, London
D	Ranasinghe	Queen Alexandra Hospital, Portsmouth	A	Stafford	Whittington Hospital, London
S	Hall	Queen Elizabeth Hospital	S	Gokani	Whittington Hospital, London
J	Wright	Queen Elizabeth Hospital	M	Charalambos	Whittington Hospital, London
S	Hall	Queen Elizabeth Hospital	A	Olajide	Whittington Hospital, London
N	Ali	Queen Elizabeth Hospital	C	Bi	Whittington Hospital, London
J	Hunt	Queen Elizabeth Hospital, Birmingham	J	Ng	Whittington Hospital, London
H	Ahmad	Queen Elizabeth Queen's Mother, Margate	S	Naeem	William Harvey Hospital, Kent
C	Ward	Queen Elizabeth Hospital, Glasgow	J	Anandarajah	Wrexham Maelor Hospital, Wales
M	Khan	Queens Medical Centre, Nottingham	A	Hill	Wythenshawe Hospital, Manchester
K	Holzman	Redhill Hospital, Surrey	C	Boulind	Yeovil District Hospital
J	Ritchie	Rotherham Hospital			
A	Hormis	Rotherham Hospital			

Ireland Trainee Emergency Research Network (I-TERN) Collaborators:

R	O'Sullivan	Bon Secours Hospital Cork, Ireland
S	Gilmartin	Children's Health Ireland at Crumlin, Ireland

S	Uí Bhroin	Children's Health Ireland at Tallaght, Ireland
P	Fitzpatrick	Children's Health Ireland at Temple Street, Ireland
A	Patton	Cork University Hospital, Ireland
M	Jee Poh Hock	Galway Hospital, Ireland
S	Graham	Mater Misericordiae University Hospital, Ireland
S	Kukaswadia	Mercy University Hospital, Ireland
C	Prendergast	Midlands Regional Hospital Tullamore, Ireland
A	Ahmed	Sligo University Hospital, Ireland
C	Dalla Vecchia	St Vincent's University Hospital, Ireland
J	Lynch	Tallaght University Hospital, Ireland
M	Grummell	Tallaght University Hospital, Ireland
I	Grossi	University Hospital Limerick, Ireland
B	MacManus	University Hospital Waterford, Ireland

Research and Audit Federation of Trainees (RAFT), Trainee Research in Intensive Care (TRIC) and Specialist Anaesthesia Trainee led Audit and Research Network (SATURN) Collaborators:

K	Samuel	North Bristol NHS Trust
A	Boyle	Royal Victoria Hospital, Belfast
A	Waite	Royal Liverpool University Hospital
B	Johnson	University of Liverpool
J	Vinagre	Children's Health Ireland at Temple Street